



Para- tt^* -bundles on the tangent bundle of an almost para-complex manifold

Lars Schaefer

► To cite this version:

Lars Schaefer. Para- tt^* -bundles on the tangent bundle of an almost para-complex manifold. 2007. hal-00143297

HAL Id: hal-00143297

<https://hal.archives-ouvertes.fr/hal-00143297>

Preprint submitted on 25 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Para- tt^* -bundles on the tangent bundle of an almost para-complex manifold

Lars Schäfer

Department Mathematik, Universität Hamburg,
Bundesstraße 55, D-20146 Hamburg, Germany
schaefer@math.uni-hamburg.de

May 24, 2006

Abstract

In this paper we study para- tt^* -bundles (TM, D, S) on the tangent bundle of an almost para-complex manifold (M, τ) . We characterise those para- tt^* -bundles with $\nabla = D + S$ induced by the one-parameter family of connections given by $\nabla^\theta = \exp(\theta\tau) \circ \nabla \circ \exp(-\theta\tau)$ and prove a uniqueness result for solutions with a para-complex connection D . Flat nearly para-Kähler manifolds and special para-complex manifolds are shown to be such solutions. We analyse which of these solutions admit metric or symplectic para- tt^* -bundles. Moreover, we give a generalisation of the notion of a para-pluriharmonic map to maps from almost para-complex manifolds (M, τ) into pseudo-Riemannian manifolds and associate to the above metric and symplectic para- tt^* -bundles generalised para-pluriharmonic maps into $\mathrm{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$, respectively into $SO_0(n, n)/U^\pi(C^n)$, where $U^\pi(C^n)$ is the para-complex analogue of the unitary group.

MSC: 53C26, 53C43, 58E20.

Keywords: Para- tt^* -geometry and para- tt^* -bundles; special para-complex and special para-Kähler manifolds; Nearly para-Kähler manifolds; para-pluriharmonic maps; pseudo-Riemannian symmetric spaces.

Contents

1	Introduction	1
2	Para-complex differential geometry	2
3	Para- tt^* -bundles	4
4	Solutions on the tangent bundle of an almost para-complex manifold	5
5	Solutions on almost para-hermitian manifolds	9
6	Para-pluriharmonic maps from almost para-complex manifolds into pseudo-Riemannian manifolds	14
7	Related para-pluriharmonic and harmonic maps	17
7.1	The classifying map of a flat nearly para-Kähler manifold	17
7.2	The dual Gauß map of a special para-Kähler manifold with torsion	19

1 Introduction

The subject of this work are para- tt^* -bundles (TM, D, S) on the tangent bundle an almost para-complex manifold (M, τ) as base. We generalise the results of the complex geometric setting which can be found in [S3]. In previous work we always supposed a para-complex base manifold. Motivated by the study of the complex case we admit non-integrable para-complex structures, since Levi-Civita flat *nearly para-Kähler manifolds* arise in this way as solutions of para- tt^* -bundles. Nearly para-Kähler manifolds as analogue of nearly Kähler manifolds were introduced in a recent paper of Ivanov and Zamkovoy [IZ]. A further class of solutions is given by **special para-Kähler manifolds** which were studied in [S2]. These geometries arise as one of the special geometries of Euclidean supersymmetry in [CMMS]. This means that para- tt^* -bundles can be seen as a common generalisation of these two geometries and give a kind of duality between them.

Now we discuss the structure of the paper. First we introduce the notion of (metric, symplectic) para- tt^* -bundles and describe these in terms of explicit geometric data. Part of this data is a family of flat connections D^θ on a vector bundle E with $\theta \in \mathbb{R}$ which is defined with help of a second connection D and a field $S \in \Gamma(T^*M \otimes \text{End}(E))$. Given an almost para-complex manifold endowed with a flat connection ∇ we can consider a natural family of flat connections defined by

$$\nabla^\theta = \exp(\theta\tau) \circ \nabla \circ \exp(-\theta\tau) \text{ for } \theta \in \mathbb{R}.$$

We analyse para- tt^* -bundles, such that $D^\theta = \nabla^{\alpha\theta}$ for some $\alpha \in \mathbb{R}$. This choice is motivated by the study of solutions coming from special para-Kähler geometry.

If we restrict to para- tt^* -bundles carrying connections D which are para-complex, i.e. which satisfy $D\tau = 0$, then these para- tt^* -bundles can be recovered uniquely from the pair (∇, τ) . Moreover, we give compatibility conditions on the pair (∇, τ) . These conditions hold for *special para-complex* and nearly para-Kähler manifolds. We discuss two classes of solutions: The first corresponds to special para-complex manifolds with torsion and the second to almost para-complex manifolds endowed with a flat connection ∇ such that (∇, τ) satisfies the nearly para-Kähler condition (with torsion). Further we study whether these para- tt^* -bundles provide *metric* or *symplectic* para- tt^* -bundles. Examples of solutions of the first type are given by special para-Kähler manifolds and solutions of the second type arise on Levi-Civita flat nearly para-Kähler manifolds. In addition we show that neither the special para-Kähler condition is compatible with symplectic para- tt^* -bundles nor the nearly para-Kähler condition is compatible with solutions of metric para- tt^* -bundles.

There exists a relation between para-pluriharmonic maps and para- tt^* -geometry which was studied in [S2]. In this work we generalise the notion of a para-pluriharmonic map to maps from almost para-complex manifolds into pseudo-Riemannian manifolds. Then we introduce \tilde{S}^1 -pluriharmonic maps which generalise S^1 -pluriharmonic maps (cf. [S3]) to para-complex geometry. This is a generalisation of the notion of associated families (see for example [ET]). We relate these \tilde{S}^1 -pluriharmonic maps to the generalisation of para-pluriharmonic maps and prove a result, which relates generalised para-pluriharmonic maps to harmonic maps. With these preparations we are able to associate para-pluriharmonic maps into $\mathrm{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$, respectively into $SO_0(n, n)/U^\pi(C^n)$, to the above metric and symplectic para- tt^* -bundles. Here $U^\pi(C^n)$ is the para-complex analogue of the unitary group.

The author is grateful to his advisor V. Cortés for the support of his work and to M.-A. Lawn-Paillusseau for related discussions.

2 Para-complex differential geometry

In this section we recall facts of para-complex differential geometry, which are needed in the later work. For more information we refer to [CMMS, IZ].

Definition 1 *A para-complex structure on a real finite dimensional vector space V is a non-trivial involution $\tau \in \mathrm{End}(V)$ such that the two eigenspaces $V^\pm := \ker(\mathrm{Id} \mp \tau)$ to the eigenvalues ± 1 of τ have the same dimension. We call the pair (V, τ) a **para-complex vector-space**.*

*An almost para-complex structure on a smooth manifold M is an endomorphism field $\tau \in \Gamma(\mathrm{End}(TM))$ such that, for all $p \in M$, τ_p is a para-complex structure on $T_p M$. It is called **integrable** if the distributions $T^\pm M = \ker(\mathrm{Id} \mp \tau)$ are integrable. An integrable almost para-complex structure on M is called a **para-complex structure on M** and a manifold M endowed with a para-complex structure is called a **para-complex manifold**.*

The integrability of a para-complex structure is obstructed (cf. [CMMS]) by the vanishing of a tensor, also called **Nijenhuis tensor**, which is defined as

$$N_\tau(X, Y) := [X, Y] + [\tau X, \tau Y] - \tau[X, \tau Y] - \tau[\tau X, Y].$$

The decomposition of the tangent bundle of an (almost) para-complex manifold M into the eigenspaces $T^\pm M$ extends to a bi-grading on the exterior algebra:

$$\Lambda^k T^* M = \bigoplus_{k=p+q} \Lambda^{p,q} T^* M \quad (2.1)$$

and induces an obvious bi-grading on exterior forms with values in a vector bundle E .

The **para-complex numbers** are the real algebra, which is generated by 1 and by the **para-complex unit** e with $e^2 = 1$. We denote the para-complex numbers by the symbol C .

For all $z = x + ey \in C$ with $x, y \in \mathbb{R}$ we define the **para-complex conjugation** $\bar{\cdot} : C \rightarrow C$, $x + ey \mapsto x - ey$ and the **real and imaginary parts** of z by

$$\Re(z) := \frac{z + \bar{z}}{2} = x, \quad \Im(z) := \frac{e(z - \bar{z})}{2} = y.$$

The free C -module C^n is a para-complex vector space where its para-complex structure is the multiplication with e and the para-complex conjugation of C extends to $\bar{\cdot} : C^n \rightarrow C^n$, $v \mapsto \bar{v}$.

Note the identity $z\bar{z} = x^2 - y^2$. Therefore the algebra C is sometimes called the **hyper-complex numbers**. The circle $\mathbb{S}^1 = \{z = x + iy \in \mathbb{C} \mid x^2 + y^2 = 1\}$ is replaced by the four hyperbolas $\{z = x + ey \in C \mid x^2 - y^2 = \pm 1\}$. $\tilde{\mathbb{S}}^1$ is defined to be the hyperbola given by the one parameter group $\{z(\theta) = \cosh(\theta) + e \sinh(\theta) \mid \theta \in \mathbb{R}\}$.

The **para-complex dimension** of a para-complex manifold M is the integer $n = \dim_C M := \frac{1}{2} \dim_{\mathbb{R}} M$.

Now we consider the **para-complexification** $TM^C = TM \otimes_{\mathbb{R}} C$ of the tangent bundle TM of an almost para-complex manifold (M, τ) and extend $\tau : TM \rightarrow TM$ C -linearly to $\tau : TM^C \rightarrow TM^C$. Then for all $p \in M$ the free C -module $T_p M^C$ decomposes as C -module into the direct sum of two free C -modules

$$T_p M^C = T_p^{1,0} M \oplus T_p^{0,1} M, \quad (2.2)$$

where $T_p^{1,0} M := \{X + e\tau X \mid X \in T_p M\}$ and $T_p^{0,1} M := \{X - e\tau X \mid X \in T_p M\}$. The subbundles $T_p^{1,0} M$ and $T_p^{0,1} M$ can be characterised as the $\pm e$ -eigenbundles of the map $\tau : TM^C \rightarrow TM^C$, i.e. $\tau = e$ on $T^{1,0} M$ and $\tau = -e$ on $T^{0,1} M$.

In the same manner we decompose $T^* M^C = \Lambda^{1,0} T^* M \oplus \Lambda^{0,1} T^* M$ into the $\pm e$ -eigenbundles of the dual para-complex structure $\tau^* : T^* M^C \rightarrow T^* M^C$. This decomposition induces a bi-grading on the C -valued exterior forms

$$\Lambda^k T^* M^C = \bigoplus_{k=p+q} \Lambda^{p,q} T^* M$$

and finally on the C -valued differential forms on M

$$\Omega_C^k(M) = \bigoplus_{k=p+q} \Omega^{p,q}(M).$$

We remark, that for the case $(1, 1)$ and $(1+, 1-)$ the two gradings induced by τ coincide in following sense: $\Lambda^{1,1} T^* M = (\Lambda^{1+,1-} T^* M) \otimes C$.

We need to introduce some additional notions.

Definition 2 Let (V, τ) be a para-complex vector space. A **para-hermitian scalar product** g on V is a pseudo-Euclidean scalar product for which τ is an anti-isometry, i.e.

$$\tau^*g = g(\tau\cdot, \tau\cdot) = -g(\cdot, \cdot).$$

A **para-hermitian vector space** (V, τ, g) is a para-complex vector space (V, τ) endowed with a para-hermitian scalar product g . The pair (τ, g) is called **para-hermitian structure** on the vector space V .

Definition 3 Let (V, τ, g) be a para-hermitian vector space. The para-unitary group of V is the automorphism group

$$U^\pi(V) := \text{Aut}(V, \tau, g) = \{L \in GL(V) \mid [L, \tau] = 0 \text{ and } L^*g = g\}.$$

Definition 4 An **almost para-hermitian manifold** (M, τ, g) is an almost para-complex manifold (M, τ) endowed with a pseudo-Riemannian metric g such that $\tau^*g = -g$. If τ is integrable, we call (M, τ, g) a **para-hermitian manifold**. The two-form $\omega := g(\tau\cdot, \cdot)$ is called the **fundamental two-form** of the almost para-hermitian manifold (M, τ, g) .

3 Para- tt^* -bundles

We generalise the notion of a para- tt^* -bundle which was introduced in [S2], by admitting a base manifold (M, τ) with a non-integrable para-complex structure τ . In this way nearly para-Kähler manifolds appear as solutions of para- tt^* -geometry on the tangent bundle TM . Further we introduce symplectic para- tt^* -bundles.

Definition 5 A **para- tt^* -bundle** or **ptt^* -bundle** (E, D, S) over an almost para-complex manifold (M, τ) is a real vector bundle $E \rightarrow M$ endowed with a connection D and a section $S \in \Gamma(T^*M \otimes \text{End } E)$ which satisfy the **ptt^* -equation**

$$R^\theta = 0 \quad \text{for all } \theta \in \mathbb{R}, \quad (3.1)$$

where R^θ is the curvature tensor of the connection D^θ defined by

$$D_X^\theta := D_X + \cosh(\theta)S_X + \sinh(\theta)S_{\tau X} \quad \text{for all } X \in TM. \quad (3.2)$$

A **metric ptt^* -bundle** (E, D, S, g) is a ptt^* -bundle (E, D, S) endowed with a possibly indefinite D -parallel fiber metric g such that for all $p \in M$

$$g(S_X Y, Z) = g(Y, S_X Z) \quad \text{for all } X, Y, Z \in T_p M. \quad (3.3)$$

A **symplectic ptt^* -bundle** (E, D, S, ω) is a ptt^* -bundle (E, D, S) endowed with the structure of a symplectic vector bundle¹ (E, ω) , such that ω is D -parallel and S is ω -symmetric, i.e. for all $p \in M$

$$\omega(S_X Y, Z) = \omega(Y, S_X Z) \quad \text{for all } X, Y, Z \in T_p M. \quad (3.4)$$

¹see D. Mc Duff and D. Salamon [McDS]

Remark 1

If (E, D, S) is a ptt^* -bundle then (E, D, S^θ) is a ptt^* -bundle for all $\theta \in \mathbb{R}$, where

$$S^\theta := D^\theta - D = \cosh(\theta)S + \sinh(\theta)S_\tau. \quad (3.5)$$

The same remark applies to metric ptt^* -bundles and symplectic ptt^* -bundles.

Like for ptt^* -bundles (E, D, S) over a para-complex manifold we find explicit equations for D and S .

Proposition 1 *Let E be a real vector bundle over an almost para-complex manifold (M, τ) endowed with a connection D and a section $S \in \Gamma(T^*M \otimes \text{End } E)$.*

Then (E, D, S) is a ptt^ -bundle if and only if D and S satisfy the following equations:*

$$R^D + S \wedge S = 0, \quad (3.6)$$

$$S \wedge S \text{ is of type } (1,1), \quad (3.7)$$

$$[D_X, S_Y] - [D_Y, S_X] - S_{[X,Y]} = 0, \quad \forall X, Y \in \Gamma(TM), \quad (3.8)$$

$$[D_X, S_{\tau Y}] - [D_Y, S_{\tau X}] - S_{\tau[X,Y]} = 0, \quad \forall X, Y \in \Gamma(TM). \quad (3.9)$$

Fixing a torsion-free connection on (M, τ) the last two equations are equivalent to

$$d^D S = 0 \quad \text{and} \quad d^D S_\tau = 0. \quad (3.10)$$

Proof: (compare [S2]) As for ptt^* -bundles over para-complex manifolds (M, τ) one calculates using the theorems of addition $2 \cosh(\theta) \sinh(\theta) = \sinh(2\theta)$, $2 \cosh^2(\theta) = 1 + \cosh(2\theta)$ and $2 \sinh^2(\theta) = \cosh(2\theta) - 1$, the (finite) decomposition of R^θ in $1, \cosh(n\theta)$ and $\sinh(n\theta)$, for $n = 1, 2$. The ptt^* -equation $R^\theta = 0$ is equivalent to the vanishing of all components. This yields the claimed equations. \square

4 Solutions on the tangent bundle of an almost para-complex manifold

In this section we consider para-complex manifolds (M, τ) endowed with a flat connection ∇ . It is natural to regard the one-parameter family of connections ∇^θ , which is defined by

$$\nabla_X^\theta Y = \exp(\theta\tau) \nabla_X (\exp(-\theta\tau) Y) \text{ for } X, Y \in \Gamma(TM), \quad (4.1)$$

where $\exp(\theta\tau) = \cosh(\theta)Id + \sinh(\theta)\tau$. The connection ∇ is flat if and only if ∇^θ is flat.

Definition 6 *Two one-parameter families of connections ∇^θ and D^θ , with $\theta \in \mathbb{R}$, on some vector bundle E are called (linear) equivalent with factor $\alpha \in \mathbb{R}$ if they satisfy the equation $\nabla^\theta = D^{\alpha\theta}$.*

In the sequel we study ptt^* -bundles (TM, D, S) such that the connection D^θ defined in equation (3.2) is linear equivalent to the connection ∇^θ defined in equation (4.1). This ansatz is motivated by our previous study of ptt^* -bundles coming from special para-complex and special para-Kähler manifolds (see [S2]).

Proposition 2 *Given an almost para-complex manifold (M, τ) with a flat connection ∇ . Then a decomposition of $\nabla = D + S$ in a connection D and a section S in $T^*M \otimes \text{End}(TM)$ defines a ptt*-bundle (TM, D, S) , such that the family of connections D^θ is linear equivalent to the family of connections ∇^θ with factor $\alpha = \pm 2$ if and only if S and D satisfy*

$$S_{\tau X} = \pm \tau S_X Y$$

and

$$-(D_X \tau)Y = \tau S_X Y + S_X \tau Y =: \{S_X, \tau\}Y$$

for all $X, Y \in \Gamma(TM)$.

Proof: One has to compute ∇^θ for $X, Y \in \Gamma(TM)$:

$$\begin{aligned} \nabla_X^\theta Y &= \exp(\theta\tau)(D_X + S_X)(\cosh(\theta)Id - \sinh(\theta)\tau)Y \\ &= D_X Y - \exp(\theta\tau) \sinh(\theta)(D_X \tau)Y \\ &\quad + (\cosh(\theta)Id + \sinh(\theta)\tau)S_X(\cosh(\theta)Id - \sinh(\theta)\tau)Y \\ &= D_X Y - (\cosh(\theta) \sinh(\theta) + \sinh^2(\theta)\tau)(D_X \tau)Y + \cosh^2(\theta)S_X Y \\ &\quad - \sinh^2(\theta)\tau S_X \tau Y - \cosh(\theta) \sinh(\theta)[S_X, \tau]Y. \end{aligned}$$

This yields with the theorems of addition, i.e.

$$2 \sinh(\theta) \cosh(\theta) = \sinh(2\theta), \quad 2 \cosh^2(\theta) = 1 + \cosh(2\theta) \quad \text{and} \quad 2 \sinh^2(\theta) = \cosh(2\theta) - 1,$$

the identity

$$\begin{aligned} \nabla_X^\theta Y &= D_X Y - \frac{1}{2} \sinh(2\theta)(D_X \tau)Y - \frac{1}{2}(\cosh(2\theta) - 1)\tau(D_X \tau)Y \\ &\quad + \frac{1}{2}(1 + \cosh(2\theta))S_X Y - \frac{1}{2}(\cosh(2\theta) - 1)\tau S_X \tau Y - \frac{1}{2} \sinh(2\theta)[S_X, \tau]Y \\ &= D_X Y + \frac{1}{2}[S_X + \tau S_X \tau + \tau D_X \tau]Y \\ &\quad + \frac{1}{2} \sinh(2\theta)[[\tau, S_X] - D_X \tau]Y \\ &\quad + \frac{1}{2} \cosh(2\theta)[S_X - \tau S_X \tau - \tau D_X \tau]Y \\ &\stackrel{!}{=} D_X Y + \cosh(\vartheta)T_X Y + \sinh(\vartheta)T_{\tau X} Y \quad \text{with } \vartheta = \pm 2\theta, \end{aligned}$$

where we have to determine $T \in \Gamma(T^*M \otimes \text{End}(TM))$.

Comparing coefficients of 1, $\cosh(2\theta)$, $\sinh(2\theta)$ gives

$$-\tau(D_X \tau)Y = S_X Y + \tau S_X \tau Y, \quad \text{or equivalently} \tag{4.2}$$

$$-(D_X \tau)Y = \tau S_X Y + S_X \tau Y = \{S_X, \tau\}Y,$$

$$T_X Y = \frac{1}{2}(S_X Y - \tau S_X \tau Y - \tau(D_X \tau)Y) \stackrel{(4.2)}{=} S_X Y, \tag{4.3}$$

$$\begin{aligned} T_{\tau X} Y &= \pm \frac{1}{2}([\tau, S_X]Y - (D_X \tau)Y) \\ &\stackrel{(4.2)}{=} \pm \frac{1}{2}(\tau S_X Y - S_X \tau Y + \tau S_X Y + S_X \tau Y) = \pm \tau S_X Y. \end{aligned} \tag{4.4}$$

The constraint on S follows from the last two equations, i.e.

$$S_{\tau X} = \pm \tau S_X Y$$

and the one on D and S from the first equation. \square

We further specialise to para-complex connections D , i.e. connections D satisfying the identity $D\tau = 0$. The existence of such connections was provided by Theorem 1 of [S2].

Corollary 1 *Let (M, τ) be an almost para-complex manifold endowed with a flat connection ∇ and let $\nabla = D + S$ be a given decomposition in a connection D and a section S in $T^*M \otimes \text{End}(TM)$, such that τ is D -parallel. Then (TM, D, S) defines a ptt*-bundle, such that the family of connections D^θ is linear equivalent to the family of connections ∇^θ with factor $\alpha = \pm 2$ if and only if S satisfies*

$$S_{\tau X} = \pm \tau S_X \text{ and } \{S_X, \tau\} = 0.$$

Proof: The second condition in proposition 2 yields using $D\tau = 0$ the equation $\{S_X, \tau\} = 0$. The first constraint of proposition 2 remains $S_{\tau X} = \pm \tau S_X \stackrel{\{S_X, \tau\}=0}{=} \mp S_X \tau$. \square

To establish a uniqueness result we need the next lemma.

Lemma 1 *Let (M, τ) be an almost para-complex manifold. Let a connection ∇ on M be given. Then the connection D and the section S of $T^*M \otimes \text{End}(TM)$ defined by*

$$S_X Y = -\frac{1}{2} \tau (\nabla_X \tau) Y \text{ and } D_X Y = \nabla_X Y - S_X Y \text{ for } X, Y \in \Gamma(TM), \quad (4.5)$$

satisfy $D\tau = 0$ and $\{S_X, \tau\} = 0$.

*Otherwise, suppose that ∇ decomposes as $\nabla = D + S$, where D is a connection on M and S is a section in $T^*M \otimes \text{End}(TM)$, such that τ is D -parallel, i.e. $D\tau = 0$ and S anti-commutes with τ , i.e. $\{S_X, \tau\} = 0$ for all $X \in \Gamma(TM)$. Then D and S are uniquely given by equation (4.5).*

Proof: We check $\nabla = D + S$ and $S_X \tau Y = -\frac{1}{2} \tau (\nabla_X \tau) \tau Y = \frac{1}{2} \tau^2 (\nabla_X \tau) Y = -\tau S_X Y$, where the second equality follows from deriving $\tau^2 = Id$. In addition it is

$$(D_X \tau) Y = (\nabla_X \tau) Y - [S_X, \tau] \stackrel{\{S_X, \tau\}=0}{=} (\nabla_X \tau) Y + 2\tau S_X = 0.$$

It rests to show the uniqueness: One starts with D' and S' having same properties and computes

$$0 = (D'_X \tau) Y = (\nabla_X \tau) Y - [S'_X, \tau] Y = (\nabla_X \tau) Y + 2\tau S'_X Y.$$

This further implies

$$S'_X Y = -\frac{1}{2} \tau (\nabla_X \tau) Y = S_X Y \text{ and } D'_X Y = \nabla_X Y - S'_X Y = \nabla_X Y - S_X Y = D_X Y.$$

\square

Corollary 1 and lemma 1 imply the following uniqueness result:

Theorem 1 *Let (M, τ) be an almost para-complex manifold endowed with a flat connection ∇ . If a decomposition of $\nabla = D + S$ in a connection D and a section S in $T^*M \otimes \text{End}(TM)$, such that τ is D -parallel, i.e. $D\tau = 0$, defines a ptt^* -bundle (TM, D, S) , such that the family of connections D^θ is linear equivalent to the family of connections ∇^θ with factor $\alpha = \pm 2$, then D and S are uniquely determined by the equations $S = -\frac{1}{2}\tau(\nabla\tau)$ and $D = \nabla - S$.*

Moreover, (TM, D, S) as above defines a ptt^ -bundle, such that the family of connections D^θ is linear equivalent to the family of connections ∇^θ with factor $\alpha = \pm 2$, if and only if τ satisfies $(\nabla_{\tau X}\tau) = \pm\tau(\nabla_X\tau)$ and D and S are given by equation (4.5).*

Some classes of examples which satisfy the condition $S_{\tau X} = \pm\tau S_X$ are given in the following propositions.

Proposition 3 *Let an almost para-complex manifold (M, τ) endowed with a connection ∇ be given and let S be the section in $T^*M \otimes \text{End}(TM)$ defined by*

$$S := -\frac{1}{2}\tau(\nabla\tau). \quad (4.6)$$

If the pair (∇, τ) satisfies one of the following conditions

- (i) (∇, τ) is special, i.e. $(\nabla_X\tau)Y = (\nabla_Y\tau)X$ for all $X, Y \in \Gamma(TM)$,*
- (ii) (∇, τ) satisfies the nearly para-Kähler condition, i.e. $(\nabla_X\tau)Y = -(\nabla_Y\tau)X$ for all $X, Y \in \Gamma(TM)$,*

then it holds $S_{\tau X}Y = -\tau S_XY$.

Proof: We prove (i) and (ii) in the same calculation:

$$(\nabla_{\tau X}\tau)Y = \pm(\nabla_Y\tau)\tau X = \mp\tau(\nabla_Y\tau)X = -\tau(\nabla_X\tau)Y.$$

This implies $S_{\tau X}Y = -\frac{1}{2}\tau(\nabla_{\tau X}\tau)Y = \frac{1}{2}\tau^2(\nabla_X\tau)Y = -\tau S_XY$.

□

Proposition 4 *Let a para-complex manifold (M, τ) endowed with a connection ∇ be given and let S be the section in $T^*M \otimes \text{End}(TM)$ defined by*

$$S := -\frac{1}{2}\tau(\nabla\tau). \quad (4.7)$$

If ∇ is (anti-)adapted, i.e. $\nabla_{\tau X} = \pm\tau\nabla_X$ for all para-holomorphic vector-fields X, Y , then it holds $S_{\tau X}Y = \pm\tau S_XY$.

Proof: Since the connection ∇ is (anti-)adapted, we obtain for all para-holomorphic vector-fields X, Y :

$$(\nabla_{\tau X}\tau)Y = \pm\tau(\nabla_X\tau)Y.$$

The following computation finishes the proof

$$S_{\tau X}Y = -\frac{1}{2}\tau(\nabla_{\tau X}\tau)Y = \mp\frac{1}{2}\tau^2(\nabla_X\tau)Y = \pm\tau S_XY.$$

□

Remark 2 *One observes that in proposition 3 the condition (i) is the symmetry of S_XY and condition (ii) is its anti-symmetry. We recall that if the connection ∇ is torsion-free, flat and special then (M, τ, ∇) is a special para-complex manifold, see [CMMS, S2]. Ptt^* -bundles coming from special para-complex manifolds and special para-Kähler manifolds were studied in [S2].*

Moreover we remark, that the second condition arises in nearly para-Kählerian geometry (compare [IZ]) and therefore is quite natural. These geometries as solutions of ptt^ -geometry are discussed later in this work.*

Finally, the notion of adapted connections appeared in the study of decompositions on (para-holomorphic) para-complex vector bundles, compare [LS].

5 Solutions on almost para-hermitian manifolds

In this section we study the question under which assumptions almost para-complex manifolds (M, τ) endowed with a flat connection ∇ such that (∇, τ) is special or satisfies the nearly para-Kähler condition define symplectic or metric ptt^* -bundles.

First we recall a lemma from tensor-algebra:

Lemma 2 *Let V be a vector-space and $\alpha \in T^3(V^*)$ an element in the third tensorial power of V^* , the dual space of V . Suppose that $\alpha(X, Y, Z)$ is symmetric (resp. anti-symmetric) in X, Y and Y, Z and $\alpha(X, Y, Z)$ is anti-symmetric (resp. symmetric) in X, Z then $\alpha = 0$.*

Proof: It is $\alpha(X, Y, Z) = \epsilon\alpha(Y, X, Z) = \epsilon\alpha(X, Z, Y)$ with $\epsilon \in \{\pm 1\}$ which implies $\alpha(X, Y, Z) = \epsilon\alpha(Y, X, Z) = \epsilon^2\alpha(Y, Z, X) = \epsilon^3\alpha(Z, Y, X)$. But further it holds $\alpha(X, Y, Z) = -\epsilon\alpha(Z, Y, X)$ and consequently $-\alpha(Z, Y, X) = \epsilon^2\alpha(Z, Y, X) = \alpha(Z, Y, X)$. This shows $\alpha = 0$. □

Proposition 5 *Let (M, τ) be an almost para-complex manifold endowed with a flat connection ∇ , such that (∇, τ) is special. Define S , a section in $T^*M \otimes \text{End}(TM)$, by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \tag{5.1}$$

then $(TM, D = \nabla - S, S)$ defines a ptt^ -bundle. If in addition (TM, D, S, ω) is a symplectic ptt^* -bundle, then it is trivial, i.e. $S = 0$.*

Proof: Due to theorem 1 and proposition 3 (TM, D, S) is a ptt^* -bundle.

Suppose, that (TM, D, S, ω) is a symplectic ptt^* -bundle. In order to prove the second part of the proposition, we define the tensor

$$\alpha(X, Y, Z) := \omega(S_X Y, Z) = g(\tau S_X Y, Z).$$

$\alpha(X, Y, Z)$ is symmetric in X, Y , since (∇, τ) is special, i.e. $\nabla \tau$ is symmetric in X, Y . Moreover, it holds

$$\begin{aligned} \alpha(X, Y, Z) &= \omega(S_X Y, Z) = -\omega(Z, S_X Y) \\ &= -\omega(Z, S_Y X) = -\omega(S_Y Z, X) = -\omega(S_Z Y, X) = -\alpha(Z, Y, X). \end{aligned}$$

This is the anti-symmetry of $\alpha(X, Y, Z)$ in X, Z . Finally we compute

$$\begin{aligned} \alpha(X, Y, Z) &= \omega(S_X Y, Z) = \omega(Y, S_X Z) \\ &= \omega(Y, S_Z X) = -\omega(S_Z X, Y) = -\alpha(Z, X, Y) = -\alpha(X, Z, Y), \end{aligned}$$

i.e. the anti-symmetry of $\alpha(X, Y, Z)$ in Y, Z .

Hence α vanishes and consequently S is zero. \square

Proposition 6 *Let an almost para-complex manifold (M, τ) endowed with a flat connection ∇ , such that (∇, τ) satisfies the nearly para-Kähler condition be given. Define S , a section S in $T^*M \otimes \text{End}(TM)$, by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \quad (5.2)$$

then $(TM, D = \nabla - S, S)$ defines a ptt^ -bundle. Suppose, that (TM, D, S, g) is a metric ptt^* -bundle, then it is trivial, i.e. $S = 0$.*

Proof: Due to theorem 1 and proposition 3 (TM, D, S) is a ptt^* -bundle.

Suppose, that it is a metric ptt^* -bundle. The proof is obtained by analysing the symmetries of the tensor

$$\alpha(X, Y, Z) := g(S_X Y, Z).$$

$\alpha(X, Y, Z)$ is anti-symmetric in X, Y , since by the nearly para-Kähler condition $\nabla \tau$ is anti-symmetric in X, Y . In addition it holds

$$\begin{aligned} \alpha(X, Y, Z) &= g(S_X Y, Z) = g(Z, S_X Y) \\ &= -g(Z, S_Y X) = -g(S_Y Z, X) = g(S_Z Y, X) = \alpha(Z, Y, X), \end{aligned}$$

which is the symmetry of $\alpha(X, Y, Z)$ in X, Z . Finally one has

$$\begin{aligned} \alpha(X, Y, Z) &= g(S_X Y, Z) = g(Y, S_X Z) \\ &= -g(Y, S_Z X) = -g(S_Z X, Y) = -\alpha(Z, X, Y) = \alpha(X, Z, Y), \end{aligned}$$

i.e. the symmetry of $\alpha(X, Y, Z)$ in Y, Z .

Hence α vanishes by the above lemma and so does S . \square

The following theorem gives solutions of symplectic para- tt^* -bundles on the tangent bundle, which are more general then the later discussed nearly para-Kähler manifolds in the sense, that we admit the connection ∇ to have torsion, but more special in the sense, that our connection ∇ has to be flat.

Theorem 2 *Let (M, τ, g) be an almost para-hermitian manifold endowed with a flat metric connection ∇ , such that the pair (∇, τ) satisfies the nearly para-Kähler condition. Define S , a section in $T^*M \otimes \text{End}(TM)$, by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \quad (5.3)$$

then $(TM, D = \nabla - S, S, \omega = g(\tau \cdot, \cdot))$ defines a symplectic ptt^ -bundle.*

The torsion T^∇ of the connection ∇ and T^D of the connection D are related by

$$T^\nabla - T^D = 2S$$

and it holds $D\tau = 0$.

Proof: Due to theorem 1 and proposition 3 (TM, D, S) is a ptt^* -bundle.

It remains to check $D\omega = 0$ and that S is ω -symmetric.

We observe, that, as g is para-hermitian and $\nabla g = 0$, $\nabla_X \tau$ is skew-symmetric with respect to g . This yields by the following calculation, that S is skew-symmetric with respect to g :

$$\begin{aligned} -2g(S_X Y, Z) &= g(\tau(\nabla_X \tau)Y, Z) = -g((\nabla_X \tau)Y, \tau Z) \\ &= g(Y, (\nabla_X \tau)\tau Z) = -g(Z, \tau(\nabla_X \tau)Y) = 2g(Y, S_X Z). \end{aligned}$$

The definition of $\omega = g(\tau \cdot, \cdot)$ and $\{S_X, \tau\} = 0$ yield the ω -symmetry of S_X . Further it holds $D = \nabla + \frac{1}{2}\tau\nabla\tau$, which implies

$$D\tau = \nabla\tau + \frac{1}{2}[\tau\nabla\tau, \tau] = 0.$$

This shows $D\omega = 0$ if and only if $Dg = 0$. But $\nabla g = 0$ and S is skew-symmetric with respect to g , so g is parallel for $D = \nabla - S$.

This shows that $(TM, D = \nabla - S, S, \omega)$ is a symplectic ptt^* -bundle.

From $\nabla = D + S$ we obtain for the torsions

$$T^\nabla(X, Y) = T^D(X, Y) + S_X Y - S_Y X = T^D(X, Y) + 2S_X Y.$$

□

We recall the definition of special para-complex and special para-Kähler manifolds (see [CMMS, S2]):

Definition 7 *A special para-Kähler manifold consists of the data (M, τ, g, ∇) where (M, τ, g) is a para-Kähler manifold with ∇ -parallel para-Kähler form and (M, τ, ∇) is a special para-complex manifold, i.e. (M, τ) is a para-complex manifold endowed with a flat and torsion-free connection ∇ such that (∇, τ) is special.*

The next theorem gives solutions of metric para- tt^* -bundles on the tangent bundle, which are more general than special para-Kähler manifolds in the sense, that we admit connections ∇ with torsion.

Theorem 3 *Let an almost para-hermitian manifold (M, τ, g) endowed with a flat connection ∇ be given. Suppose that (∇, τ) is special and the two-form $\omega = g(\tau \cdot, \cdot)$ is ∇ -parallel. Define S , a section in $T^*M \otimes \text{End}(TM)$, by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \quad (5.4)$$

then $(TM, D = \nabla - S, S, g)$ defines a metric ptt^ -bundle. Moreover, the connections D and ∇ have the same torsion and we have $D\tau = 0$.*

If in addition ∇ is torsion-free, then D is the Levi-Civita connection of g , (M, τ, g) is a para-Kähler manifold and (M, τ, g, ∇) is a special para-Kähler manifold.

Proof: Due to theorem 1 and proposition 3 (TM, D, S) defines a ptt^* -bundle.

It remains to prove $Dg = 0$ and that S is g -symmetric.

First we remark that $\omega(\tau X, Y) = -\omega(X, \tau Y)$ as g is para-hermitian. This implies by $\nabla\omega = 0$ the ω -skew-symmetry of $\nabla_X\tau$, which yields that $S_X = -\frac{1}{2}\tau(\nabla_X\tau)$ is ω -skew-symmetric, since $\tau(\nabla_X\tau) = -(\nabla_X\tau)\tau$. Finally $\{S_X, \tau\} = 0$ shows the g -symmetry of S_X . Moreover, we compute

$$D\tau = \nabla\tau + \frac{1}{2}[\tau\nabla\tau, \tau] = 0$$

and consequently $Dg = 0$ is equivalent to $D\omega = 0$.

From $\nabla\omega = 0$ and the ω -skew-symmetry of S it follows $D\omega = (\nabla - S)\omega = 0$.

From the symmetry of $\nabla\tau$, i.e. $(\nabla_X\tau)Y = (\nabla_Y\tau)X$ for all $X, Y \in TM$ we obtain $S_XY = S_YX$. This shows, that the torsions of the connections ∇ and D coincide.

Suppose now that ∇ is torsion-free. This implies, that $D = \nabla - S$ is torsion-free and consequently the Levi-Civita-connection of g . Further the equation $\nabla\omega = 0$ implies $d\omega = 0$, since ∇ is torsion-free. As $D\tau = 0$ and D is torsion-free, τ is integrable. Hence (M, τ, g) is para-Kähler. In addition (M, τ, ∇) is special para-complex by the conditions on ∇ and τ . Summarising (M, τ, g, ∇) is a special para-Kähler manifold. \square

In [S2] we treated special para-Kähler solutions of ptt^* -geometry in more details.

Now we are going to apply the above results to nearly para-Kähler manifolds. In order to do this we recall the notion of a nearly para-Kähler manifold which was recently introduced by Ivanov and Zamkovoy [IZ].

Definition 8 *An almost para-hermitian manifold (M, τ, g) is called nearly para-Kähler manifold, if its Levi-Civita connection $\nabla = \nabla^g$ satisfies the equation*

$$(\nabla_X\tau)Y = -(\nabla_Y\tau)X, \quad \forall X, Y \in \Gamma(TM). \quad (5.5)$$

A nearly para-Kähler manifold is called strict, if $\nabla\tau \neq 0$.

We recall that the tensor $\nabla\tau$ defines two tensors A and B

$$A(X, Y, Z) := g((\nabla_X\tau)Y, Z) \text{ and } B(X, Y, Z) := g((\nabla_X\tau)Y, \tau Z) \text{ with } X, Y, Z \in TM,$$

which are both (real) three-forms of type $(3, 0) + (0, 3)$.

A connection of particular importance in nearly para-Kähler geometry is the connection $\bar{\nabla}$ defined by

$$\bar{\nabla}_X Y := \nabla_X Y - \frac{1}{2}(\nabla_X \tau)\tau Y, \text{ for all } X, Y \in \Gamma(TM). \quad (5.6)$$

We remark, that $\bar{\nabla}$ is the unique connection with totally skew-symmetric torsion satisfying $\bar{\nabla}g = 0$ and $\bar{\nabla}\tau = 0$ (compare [IZ]).

The torsion of the connection $\bar{\nabla}$ is given by

$$T^{\bar{\nabla}}(X, Y) = -(\nabla_X \tau)\tau Y, \text{ for all } X, Y \in \Gamma(TM) \quad (5.7)$$

and it vanishes if and only if (M, τ, g) is a para-Kähler manifold.

Corollary 2 *Let (M, τ, g) be a nearly para-Kähler manifold, such that its Levi-Civita connection ∇ is flat and let S be the section in $T^*M \otimes \text{End}(TM)$ defined by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \quad (5.8)$$

then $(TM, \bar{\nabla}, S)$ defines a ptt^ -bundle. Suppose, that $(TM, \bar{\nabla}, S, g)$ is a metric ptt^* -bundle, then it is trivial, i.e. $S = 0$ and consequently (M, τ, g) is para-Kähler.*

Proof: If one puts $D = \bar{\nabla}$ we are in the situation of proposition 6. □

Remark 3 *In common work with V. Cortés [CS2] we gave a constructive classification of flat nearly pseudo-Kähler manifolds. The application of the methods of [CS2] to nearly para-Kähler manifolds is work in progress [CS3].*

Theorem 4 *Let (M, τ, g) be a nearly para-Kähler manifold, such that its Levi-Civita connection ∇ is flat. Let S be the section in $T^*M \otimes \text{End}(TM)$ defined by*

$$S := -\frac{1}{2}\tau(\nabla\tau), \quad (5.9)$$

then $(TM, \bar{\nabla}, S, \omega := g(\tau\cdot, \cdot))$ is a symplectic ptt^ -bundle. Further it holds*

$$B(X, Y, Z) = 2g(S_X Y, Z) \text{ and } \bar{\nabla}\tau = 0. \quad (5.10)$$

Proof: By setting $D = \bar{\nabla}$ we are in the situation of theorem 2. In addition it holds

$$2g(S_X Y, Z) = -g(\tau(\nabla_X \tau)Y, Z) = g((\nabla_X \tau)Y, \tau Z) = B(X, Y, Z).$$

□

6 Para-pluriharmonic maps from almost para-complex manifolds into pseudo-Riemannian manifolds

In this section we generalise the notion of a para-pluriharmonic map to maps from almost para-complex manifolds to pseudo-Riemannian manifolds and we introduce the para-complex analogue of an associated family (compare [ET] for the complex setting). Afterwards we show that maps admitting the para-complex analogue of an associated family give rise to a para-pluriharmonic map and we give conditions under which a para-pluriharmonic map is harmonic.

Let (M, τ) be an almost para-complex manifold of real dimension $2n$. In complex geometry it is well-known (compare [KN]) that on every almost complex manifold (M, J) there exists a connection with torsion $T = \frac{1}{4}N_J$ where N_J is the Nijenhuis tensor of the complex structure J . This result was generalised to para-complex geometry in Theorem 1 of [S2], which states that on an almost para-complex manifold (M, τ) there exists a para-complex connection with torsion $T = -\frac{1}{4}N_\tau$ where N_τ is the Nijenhuis tensor of the para-complex structure τ .

Definition 9 *Let (M, τ) be an almost para-complex manifold. A connection D on the tangent bundle of M is called **nice** if it is para-complex and its torsion T satisfies $T = \lambda N_\tau$ with a smooth function λ on M .*

We introduce the notion of a para-pluriharmonic map from an almost para-complex manifold:

Definition 10 *Let (M, τ, D) be an almost para-complex manifold endowed with a nice connection D on TM and N be a smooth manifold endowed with a connection ∇^N . Denote by ∇ the connection on $T^*M \otimes f^*TN$ which is induced by D and ∇^N .*

*A smooth map $f : M \rightarrow N$ is called **para-pluriharmonic** if and only if it satisfies the equation*

$$(\nabla df)^{1,1} = 0. \quad (6.1)$$

Remark 4 *First we remark, that for a para-complex manifold (M, τ) and a pseudo-Riemannian target manifold (N, h) with its Levi-Civita connection ∇^h the para-pluriharmonic equation (6.1) does not depend on the connection D if D is chosen in an appropriate class (compare [S2]). In fact nice connections on para-complex manifolds belong to this class. A famous case are para-Kähler manifolds (M, τ, g) , where D is taken to be the Levi-Civita connection.*

To deal with associated families of para-pluriharmonic maps we need to recall an integrability condition satisfied by the differential of a smooth map. Denote by N a smooth manifold with a connection ∇^N on its tangent bundle having torsion tensor T^N . Let a second smooth manifold M and a smooth map $f : M \rightarrow N$ be given. The differential $F := df : TM \rightarrow f^*TN = E$ of the map f induces a vector bundle homomorphism

between the tangent bundle TM of M and the pull-back of TN via f . The bundle homomorphism $T^E : \Lambda^2 E \rightarrow E$ induced by the torsion T^N of N satisfies the identity

$$\nabla_V^E F(W) - \nabla_W^E F(V) - F([V, W]) = T^E(F(V), F(W)), \quad (6.2)$$

where $\nabla^E = f^* \nabla^N$ denotes the pull-back connection, i.e. the connection which is induced on E by ∇^N and where $V, W \in \Gamma(TM)$.

In the rest of this section we denote by D a nice connection on the almost para-complex manifold (M, τ) . Under this assumption we restate the condition (6.2):

$$\begin{aligned} T^E(F(V), F(W)) &= \nabla_V^E F(W) - \nabla_W^E F(V) - F([V, W]) \\ &= \nabla_V^E F(W) - \nabla_W^E F(V) \\ &\quad - F(D_V W) + F(D_W V) + F(T(V, W)) \\ &= \nabla_V^E F(W) - \nabla_W^E F(V) \\ &\quad - F(D_V W) + F(D_W V) + \lambda F(N_\tau(V, W)) \\ &= (\nabla_V F)W - (\nabla_W F)V + \lambda F(N_\tau(V, W)), \end{aligned} \quad (6.3)$$

where ∇ is the connection induced on $T^*M \otimes E$ by D and ∇^E .

Later in this work we consider the case where N is a pseudo-Riemannian symmetric space with its Levi-Civita connection ∇^N .

Let $\alpha \in \mathbb{R}$ and define $\mathcal{R}_\alpha : TM \rightarrow TM$ as

$$\mathcal{R}_\alpha(X) = \cosh(\alpha)X + \sinh(\alpha)\tau X.$$

This is a parallel endomorphism field on the tangent bundle TM of M . The eigenvalues of which are seen to be e^{α} on $T^{1,0}M$ and $e^{-\alpha}$ on $T^{0,1}M$.

An associated family for f is a family of maps $f_\alpha : M \rightarrow N$ with $\alpha \in \mathbb{R}$ and $f_0 = f$, such that

$$\Phi_\alpha \circ df_\alpha = df \circ \mathcal{R}_\alpha, \quad \forall \alpha \in \mathbb{R}, \quad (6.4)$$

for some bundle isomorphism $\Phi_\alpha : f_\alpha^* TN \rightarrow f^* TN$, $\alpha \in \mathbb{R}$, which is parallel with respect to ∇^N , i.e. Φ_α satisfies

$$\Phi_\alpha \circ (f_\alpha^* \nabla^N) = (f^* \nabla^N) \circ \Phi_\alpha.$$

One can check, that each map f_α of an associated family itself admits an associated family.

Theorem 5 *Let (M, τ) be an almost para-complex manifold endowed with a nice connection D , N be a smooth manifold endowed with a torsion-free connection ∇^N and $f : (M, D, \tau) \rightarrow (N, \nabla^N)$ be a smooth map admitting an associated family f_α , then f is para-pluriharmonic. More precisely, each map of the associated family f_α is para-pluriharmonic.*

Proof: Since Φ_α is parallel with respect to ∇^N , ∇^N is torsion-free and D is nice, we are able to apply equation (6.3) to the family $df_\alpha = F_\alpha = \Phi_\alpha^{-1} \circ df \circ \mathcal{R}_\alpha$ and find

$$(\nabla_V F_\alpha)W - (\nabla_W F_\alpha)V + \lambda F_\alpha(N_\tau(V, W)) = 0.$$

As \mathcal{R}_α is D -parallel we obtain

$$(\nabla_X F_\alpha) = \Phi_\alpha^{-1} \circ (\nabla_X F) \circ \mathcal{R}_\alpha.$$

If $Z = X - e\tau X$ and $W = Y + e\tau Y$ have different type it holds $N_\tau(Z, W) = 0$, where we extended the Nijenhuis tensor para-complex linearly. This yields

$$(\nabla_V F_\alpha)W = (\nabla_W F_\alpha)V, \quad \forall \alpha \in \mathbb{R}$$

and further we get

$$\begin{aligned} (\nabla_Z F_\alpha)W &= e^{e\alpha} \Phi_\alpha^{-1}(\nabla_Z F)W, \\ (\nabla_W F_\alpha)Z &= e^{-e\alpha} \Phi_\alpha^{-1}(\nabla_W F)Z = e^{-e\alpha} \Phi_\alpha^{-1}(\nabla_Z F)W, \end{aligned}$$

for all $\alpha \in \mathbb{R}$. Since this should coincide, it follows $(\nabla df)^{(1,1)} = 0$, i.e. $f : (M, D, \tau) \rightarrow (N, \nabla^N)$ is para-pluriharmonic. The rest follows, since each map of the associated family f_α admits an associated family $g_\beta = f_{(\alpha+\beta)}$. \square

This is the motivation of the following definition:

Definition 11 *Let (M, τ) be an almost para-complex manifold endowed with a nice connection D , N be a smooth manifold endowed with a torsion-free connection ∇^N . A smooth map $f : (M, D, \tau) \rightarrow (N, \nabla^N)$ is said to be \tilde{S}^1 -pluriharmonic if and only if it admits an associated family.*

Given a para-hermitian metric g on M then in general a nice connection D is not the Levi-Civita connection ∇^g of g . Therefore the para-pluriharmonic equation (6.1) does not imply the harmonicity of f . If in addition the tensor $D - \nabla^g$ is trace-free the para-pluriharmonic equation implies the harmonic equation. This is true in the case of a special para-Kähler manifold (M, τ, g, ∇) and for a nearly para-Kähler manifold, where $D = \bar{\nabla}$ and $\bar{\nabla} - \nabla^g$ is skew-symmetric.

Proposition 7 *Let (M, τ, g) be an almost para-hermitian manifold endowed with a nice connection D , N be a pseudo-Riemannian manifold with its Levi-Civita connection ∇^N . Suppose, that the tensor $S = \nabla^g - D$ is trace-free. Then a para-pluriharmonic map $f : M \rightarrow N$ is harmonic.*

Proof: We calculate the expression

$$\begin{aligned} \text{tr}_g(\nabla df) &= \sum_i g(e_i, e_i) [\nabla_{e_i}^E df(e_i) - df(D_{e_i} e_i)] \\ &= \sum_i g(e_i, e_i) [\nabla_{e_i}^E df(e_i) - df((\nabla^g - S)_{e_i} e_i)] \\ &= \sum_i g(e_i, e_i) [\nabla_{e_i}^E df(e_i) - df(\nabla_{e_i}^g e_i)] \\ &= \text{tr}_g(\tilde{\nabla}^g df) \end{aligned}$$

where $\tilde{\nabla}^g$ is the connection induced on $T^*M \otimes E$ by ∇^g and ∇^E and e_i is an orthogonal basis for g on TM . Since g is para-hermitian, we obtain from the para-pluriharmonic equation

$$\text{tr}_g(\nabla df) = \text{tr}_g(\nabla df^{(1,1)}) = 0.$$

\square

7 Related para-pluriharmonic and harmonic maps

7.1 The classifying map of a flat nearly para-Kähler manifold

In this section we consider simply connected almost para-hermitian manifolds (M, τ, g) endowed with a flat metric connection ∇ such that (∇, τ) satisfies the nearly para-Kähler condition.

In particular, simply connected flat nearly para-Kähler manifolds (M^{2n}, τ, g) , i.e. nearly para-Kähler manifolds (M, τ, g) with flat Levi-Civita connection ∇^g are of this type.

Since (M, g, ∇) is simply connected and flat, we may identify by fixing a ∇ -parallel frame s_0 its tangent bundle TM with $(M \times V, \langle \cdot, \cdot \rangle)$, where $V = C^n = (\mathbb{R}^{2n}, j_0)$ is endowed with the standard scalar product $\langle \cdot, \cdot \rangle$ of signature (n, n) on $V = \mathbb{R}^{2n}$.

The compatible para-complex structure τ defines via this identification a map

$$\tau : M \rightarrow \mathcal{P}(V, \langle \cdot, \cdot \rangle),$$

where $\mathcal{P}(V, \langle \cdot, \cdot \rangle)$ is the set of para-complex structures on V which are compatible with $\langle \cdot, \cdot \rangle$ and the orientation.

One can consider $\mathcal{P}(V, \langle \cdot, \cdot \rangle)$ as a subset in the vector space $\mathfrak{so}(n, n) = \mathfrak{so}(V) \subset \text{Mat}(\mathbb{R}^{2n})$ characterised by the equation

$$f(j) = \mathbb{1}_{2n}, \quad (7.1)$$

where $f : \text{Mat}(\mathbb{R}^{2n}) \rightarrow \text{Mat}(\mathbb{R}^{2n})$ is given by $f : A \mapsto A^2$. The differential of this map is $df_A(H) = \{A, H\}$ for $A, H \in \text{Mat}(\mathbb{R}^{2n})$. We remark, that elements satisfying the equation (7.1) define automatically para-complex structures, since they are trace-free and hence their eigenspaces to the eigenvalues ± 1 have the same dimension. The differential df has constant rank in points j satisfying the equation (7.1), since one sees

$$\begin{aligned} \ker df_j &= \{A \in \mathfrak{so}(V) \mid \{j, A\} = 0\}, \\ \text{im } df_j &\cong \{A \in \mathfrak{so}(V) \mid [j, A] = 0\} \cong \mathfrak{u}^\pi(C^n). \end{aligned}$$

Applying the regular value theorem $\mathcal{P}(V, \langle \cdot, \cdot \rangle)$ is shown to be a submanifold of $\mathfrak{so}(V)$. Its tangent space at $j \in \mathcal{P}(V, \langle \cdot, \cdot \rangle)$ is

$$T_j \mathcal{P}(V, \langle \cdot, \cdot \rangle) = \ker df_j = \{A \in \mathfrak{so}(V) \mid \{j, A\} = 0\}. \quad (7.2)$$

Moreover, $\mathcal{P}(V, \langle \cdot, \cdot \rangle)$ can be identified with the pseudo-Riemannian symmetric space $SO_0(n, n)/U^\pi(C^n)$, where $SO_0(n, n)$ is the identity component of the special pseudo-orthogonal group $SO(n, n)$ and $U^\pi(C^n)$ is the para-unitary group, by the map

$$\begin{aligned} \Phi &: SO_0(n, n)/U^\pi(C^n) \rightarrow \mathcal{P}(V, \langle \cdot, \cdot \rangle), \\ gK &\mapsto g j_0 g^{-1}, \end{aligned}$$

which maps the canonical base point $o = eK$ to j_0 .

An element $j \in \mathcal{P}(V, \langle \cdot, \cdot \rangle)$ defines a symmetric decomposition of $\mathfrak{so}(V)$ by

$$\begin{aligned} \mathfrak{p}(j) &= \{A \in \mathfrak{so}(V) \mid \{j, A\} = 0\}, \\ \mathfrak{k}(j) &= \{A \in \mathfrak{so}(V) \mid [j, A] = 0\} \cong \mathfrak{u}^\pi(C^n). \end{aligned}$$

In particular we note $\mathfrak{k}(j_0) = \mathfrak{u}^\pi(C^n)$. Moreover, one observes $T_j \mathcal{P}(V, \langle \cdot, \cdot \rangle) = \mathfrak{p}(j)$. Let $\tilde{j} \in SO_0(n, n)/U^\pi(C^n)$ and $j = \Phi(\tilde{j})$, then $T_{\tilde{j}} SO_0(n, n)/U^\pi(C^n)$ is canonically identified with $\mathfrak{p}(j)$. We determine now the differential of the above identification:

Proposition 8 *Let $\Psi = \Phi^{-1} : \mathcal{P}(V, \langle \cdot, \cdot \rangle) \rightarrow SO_0(n, n)/U^\pi(C^n)$. Then it holds at $j \in \mathcal{P}(V, \langle \cdot, \cdot \rangle)$*

$$d\Psi : T_j \mathcal{P}(V, \langle \cdot, \cdot \rangle) \ni X \mapsto -\frac{1}{2}j^{-1}X \in \mathfrak{p}(j). \quad (7.3)$$

The next theorem constructs the desired para-pluriharmonic map.

Theorem 6 *Let (M, τ, g) be a simply connected almost para-hermitian manifold endowed with a flat metric connection ∇ such that (∇, τ) satisfies the nearly para-Kähler condition, then $(TM, D = \nabla - S, S = -\frac{1}{2}\tau(\nabla\tau), \omega = g(\tau\cdot, \cdot))$ defines a symplectic ptt*-bundle and the matrix of τ in a D^θ -flat frame $s^\theta = (s_i^\theta)$ defines an \tilde{S}^1 -pluriharmonic map $\tilde{\tau}^\theta : M \rightarrow \mathcal{P}(V, \langle \cdot, \cdot \rangle) \rightarrow SO_0(n, n)/U^\pi(C^n)$.*

In particular, given a nice connection D on M the map

$$\tilde{\tau}^\theta : (M, \tau, D) \rightarrow SO_0(n, n)/U^\pi(C^n)$$

is para-pluriharmonic.

Proof: One observes $D^\theta g = 0$, as one has $\nabla g = 0$ and $S_X^\theta := \cosh(\theta)S_X + \sinh(\theta)S_{\tau X}$ takes values in $\mathfrak{so}(V)$. This means we can choose for each θ the D^θ -flat frame s^θ orthonormal, such that $s^{\theta=0} = s_0$. Using $D\tau = 0$ (compare theorem 1 and lemma 1) we obtain

$$Xg(\tau s_i^\theta, s_j^\theta) = g(D_X^\theta(\tau s_i^\theta), s_j^\theta) = g((D_X^\theta \tau)s_i^\theta, s_j^\theta) = g([S_X^\theta, \tau]s_i^\theta, s_j^\theta) = -2g(\tau S_X^\theta s_i^\theta, s_j^\theta).$$

Let S^{s^θ} and τ^{s^θ} be the representation of S and τ in the frame s^θ , then we have

$$(\tau^{s^\theta})^{-1}X(\tau^{s^\theta}) = -2S^{s^\theta}$$

or

$$d\tilde{\tau}^\theta = (s^\theta)^{-1} \circ S^\theta \circ s^\theta.$$

Here the frame s^θ is seen as a map $s^\theta : M \times V \rightarrow TM$. For $X \in \Gamma(TM)$ we obtain

$$\begin{aligned} d\tilde{\tau}^\theta(X) &= (s^\theta)^{-1} \circ S_X^\theta \circ (s^\theta) = (s^\theta)^{-1} \circ S_{\mathcal{R}_\theta X} \circ (s^\theta) \\ &= ((s^\theta)^{-1} s^0) \circ d\tilde{\tau}(\mathcal{R}_\theta X) \circ ((s^0)^{-1} s^\theta) \\ &= Ad_{\alpha_\theta}^{-1} \circ d\tilde{\tau}(\mathcal{R}_\theta X) = \Phi_\theta^{-1} \circ d\tilde{\tau}(\mathcal{R}_\theta X), \end{aligned}$$

where $\alpha_\theta = (s^\theta)^{-1} s^0$ is the frame change from s_0 to s_θ and $\Phi_\theta = Ad_{\alpha_\theta}$. Φ_θ is parallel with respect to the Levi-Civita connection on $SO_0(n, n)/U^\pi(C^n)$ and hence $\tilde{\tau}^\theta$ is \tilde{S}^1 -pluriharmonic. Given a nice connection D on M theorem 5 shows that $\tilde{\tau}^\theta$ is para-pluriharmonic. \square

The next corollary emphasises the nearly para-Kähler setting:

Corollary 3 *Let (M, τ, g) be a simply connected flat nearly para-Kähler manifold and $(TM, \bar{\nabla} = \nabla^g - S, S = -\frac{1}{2}\tau(\nabla\tau), \omega(\cdot, \cdot) = g(\tau\cdot, \cdot))$ the associated symplectic ptt*-bundle, then the matrix of τ in a D^θ -flat frame $s^\theta = (s_i^\theta)$ defines an \tilde{S}^1 -pluriharmonic map $\tilde{\tau}^\theta : M \rightarrow \mathcal{P}(V, \langle \cdot, \cdot \rangle) \rightarrow SO_0(n, n)/U^\pi(C^n)$.*

For nearly para-Kähler manifolds one can get more precise information about the map $\tilde{\tau}^\theta$:

Theorem 7 *Let (M, τ, g) be a simply connected flat nearly para-Kähler manifold and $(TM, \bar{\nabla} = \nabla^g - S, S = -\frac{1}{2}\tau(\nabla\tau), \omega(\cdot, \cdot) = g(\tau\cdot, \cdot))$ the associated symplectic ptt*-bundle. Then the connection $\bar{\nabla}$ is nice and the matrix of τ in a D^θ -flat frame $s^\theta = (s_i^\theta)$ defines a para-pluriharmonic map $\tilde{\tau}^\theta : (M, \tau, \bar{\nabla}) \rightarrow \mathcal{P}(V, \langle \cdot, \cdot \rangle) \rightarrow SO_0(n, n)/U^\pi(C^n)$. Moreover, the map $\tilde{\tau}^\theta$ is harmonic.*

Proof: We have to prove, that $\bar{\nabla}$ is nice. Therefore we rewrite the Nijenhuis tensor

$$\begin{aligned} N_\tau(X, Y) &= (\nabla_{\tau X}\tau)Y - (\nabla_{\tau Y}\tau)X - \tau(\nabla_X\tau)Y + \tau(\nabla_Y\tau)X \\ &= -4\tau(\nabla_X\tau)Y, \end{aligned}$$

where the second equality follows from the nearly para-Kähler condition and by

$$(\nabla_{\tau X}\tau)Y = -(\nabla_Y\tau)\tau X = \tau(\nabla_Y\tau)X = -\tau(\nabla_X\tau)Y.$$

On the other hand the torsion of $\bar{\nabla}$ was given in equation (5.7) by

$$T^{\bar{\nabla}}(X, Y) = \tau(\nabla_X\tau)Y$$

and consequently $\bar{\nabla}$ is nice.

Due to corollary 3 the map $\tilde{\tau}^\theta$ is \tilde{S}^1 -pluriharmonic. Since $\bar{\nabla}$ is nice, theorem 5 implies that $\tilde{\tau}^\theta$ is para-pluriharmonic. From the skew-symmetry of S and proposition 7 we obtain that $\tilde{\tau}^\theta$ is harmonic. \square

7.2 The dual Gauß map of a special para-Kähler manifold with torsion

In this subsection we consider a simply connected almost para-hermitian manifold (M, τ, g) with a flat connection ∇ , such that (∇, τ) is special and the two-form $\omega = g(\tau\cdot, \cdot)$ is ∇ -parallel.

Using the flat connection ∇ we identify by fixing a ∇ -parallel symplectic frame s_0 the tangent bundle (TM, ω) with $(M \times V, \omega_0)$ where $V = \mathbb{R}^{2n}$ and ω_0 is its standard symplectic form.

The compatible para-complex structure τ is seen as a map

$$\tau : M \rightarrow \mathcal{P}(V, \omega_0),$$

where $\mathcal{P}(V, \omega_0)$ is the set of para-complex structures on V which are compatible with ω_0 . Now we discuss the differential geometry of $\mathcal{P}(V, \omega_0)$, where ω_0 is the standard symplectic

form on $V = C^n = (\mathbb{R}^{2n}, j_0)$.

First, we consider $\mathcal{P}(V, \omega_0)$ as a subset of the vector space $\mathfrak{sp}(\mathbb{R}^{2n}) \subset \text{Mat}(\mathbb{R}^{2n})$ characterised by the set of equations

$$f(j) = \mathbb{1}_{2n}, \quad (7.4)$$

where $f : \text{Mat}(\mathbb{R}^{2n}) \rightarrow \text{Mat}(\mathbb{R}^{2n})$ is given as in the last section. Again, df has constant rank in points j satisfying the equation (7.4), since one sees

$$\begin{aligned} \ker df_j &= \{A \in \mathfrak{sp}(\mathbb{R}^{2n}) \mid \{j, A\} = 0\}, \\ \text{im } df_j &\cong \{A \in \mathfrak{sp}(\mathbb{R}^{2n}) \mid [j, A] = 0\} \cong \mathfrak{u}^\pi(C^r). \end{aligned}$$

Applying the regular value theorem we obtain that $\mathcal{P}(V, \omega_0)$ is a submanifold of $\mathfrak{sp}(\mathbb{R}^{2n})$. Its tangent space at $j \in \mathcal{P}(V, \omega_0)$ is

$$T_j \mathcal{P}(V, \omega_0) = \ker df_j = \{A \in \mathfrak{sp}(\mathbb{R}^{2n}) \mid \{j, A\} = 0\}. \quad (7.5)$$

In addition the manifold $\mathcal{P}(V, \omega_0)$ can be identified with the pseudo-Riemannian symmetric space $\text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$ by the map

$$\begin{aligned} \Phi : \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n) &\rightarrow \mathcal{P}(V, \omega_0), \\ gK &\mapsto g j_0 g^{-1}, \end{aligned}$$

which maps the canonical base point $o = eK$ to j_0 .

Any $j \in \mathcal{P}(V, \omega_0)$ defines a symmetric decomposition of $\mathfrak{sp}(\mathbb{R}^{2n})$ by

$$\begin{aligned} \mathfrak{p}(j) &= \{A \in \mathfrak{sp}(\mathbb{R}^{2n}) \mid \{j, A\} = 0\}, \\ \mathfrak{k}(j) &= \{A \in \mathfrak{sp}(\mathbb{R}^{2n}) \mid [j, A] = 0\} \cong \mathfrak{u}^\pi(C^r). \end{aligned}$$

In particular it is $\mathfrak{k}(j_0) = \mathfrak{u}^\pi(C^r)$. Moreover, one observes $T_j \mathcal{P}(V, \omega_0) = \mathfrak{p}(j)$.

Let $\tilde{j} \in \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$ and $j = \Phi(\tilde{j})$, then $T_{\tilde{j}} \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$ is canonically identified with $\mathfrak{p}(j)$ and for the differential of the identification we have:

Proposition 9 *Let $\Psi = \Phi^{-1} : \mathcal{P}(V, \omega_0) \rightarrow \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$. Then it holds at $j \in \mathcal{P}(V, \omega_0)$*

$$d\Psi : T_j \mathcal{P}(V, \omega_0) \ni X \mapsto -\frac{1}{2}j^{-1}X \in \mathfrak{p}(j). \quad (7.6)$$

Recall, that under the above assumptions $(TM, D = \nabla - S, S = -\frac{1}{2}\tau(\nabla\tau), g)$ defines a metric ptt^* -bundle. Analogue to the last section we obtain:

Theorem 8 *Let (M, τ, g) be a simply connected almost para-hermitian manifold endowed with a flat connection ∇ , such that (∇, τ) is special and the two-form $\omega = g(\tau \cdot, \cdot)$ is ∇ -parallel and let $(TM, D = \nabla - S, S = -\frac{1}{2}\tau(\nabla\tau), g)$ be the associated metric ptt^* -bundle. Then the matrix of τ in a D^θ -flat frame $s^\theta = (s_i^\theta)$ defines an \tilde{S}^1 -pluriharmonic map $\tilde{\tau}^\theta : M \rightarrow \mathcal{P}(V, \omega_0) \rightarrow \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$.*

In particular, given a nice connection D on (M, τ) then the map $\tilde{\tau}^\theta : (M, \tau, D) \rightarrow \text{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$ is para-pluriharmonic.

Proof: Since we have $D^0\omega = \nabla\omega = (D + S)\omega = 0$ and $S_X^\theta := \cosh(\theta)S_X + \sinh(\theta)S_{\tau X}$ is skew-symmetric with respect to ω , we obtain $D\omega = 0$ and $D^\theta\omega = 0$. Hence one can choose for each θ the D^θ -parallel frame s^θ as a symplectic frame, such that $s^{\theta=0} = s_0$. This yields using $D\tau = 0$ (cf. theorem 1 and lemma 1)

$$X.\omega(\tau s_i^\theta, s_j^\theta) = \omega(D_X^\theta(\tau s_i^\theta), s_j^\theta) = \omega((D_X^\theta\tau)s_i^\theta, s_j^\theta) = \omega([S_X^\theta, \tau]s_i^\theta, s_j^\theta) = -2\omega(\tau S_X^\theta s_i^\theta, s_j^\theta).$$

Denote by $S^{s^\theta}, \tau^{s^\theta}$ the representation of S and τ in the frame s^θ , then we get

$$(\tau^{s^\theta})^{-1}X(\tau^{s^\theta}) = -2S^{s^\theta}$$

or

$$d\tilde{\tau}^\theta = (s^\theta)^{-1} \circ S^\theta \circ s^\theta,$$

where the frame s^θ is seen as a map $s^\theta : M \times V \rightarrow TM$. This shows for $X \in \Gamma(TM)$

$$\begin{aligned} d\tilde{\tau}^\theta(X) &= (s^\theta)^{-1} \circ S_X^\theta \circ (s^\theta) = (s^\theta)^{-1} \circ S_{\mathcal{R}_\theta X} \circ (s^\theta) \\ &= ((s^\theta)^{-1}s^0) \circ d\tilde{\tau}(\mathcal{R}_\theta X) \circ ((s^0)^{-1}s^\theta) \\ &= Ad_{\alpha_\theta}^{-1} \circ d\tilde{\tau}(\mathcal{R}_\theta X) = \Phi_\theta^{-1} \circ d\tilde{\tau}(\mathcal{R}_\theta X), \end{aligned}$$

where $\alpha_\theta = (s^\theta)^{-1}s^0$ denotes the frame change from s_0 to s_θ and $\Phi_\theta = Ad_{\alpha_\theta}$ which is parallel with respect to the Levi-Civita connection on $\mathrm{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$. In other words we have found an associated family. Given a nice connection D on (M, τ) theorem 5 shows that $\tilde{\tau}^\theta$ is para-pluriharmonic. \square

In the case where the above ptt^* -bundle comes from a special para-Kähler manifold we have the theorem:

Theorem 9 *Let (M, τ, g, ∇) be a simply connected special para-Kähler manifold and $(TM, D = \nabla - S, S = -\frac{1}{2}\tau\nabla\tau, g)$ be the associated metric ptt^* -bundle, then the matrix of τ in a D^θ -flat frame $s^\theta = (s_i^\theta)$ defines a para-pluriharmonic map $\tilde{\tau}^\theta : (M, \tau, D) \rightarrow \mathrm{Sp}(\mathbb{R}^{2n})/U^\pi(C^n)$. Further $\tilde{\tau}^\theta$ is harmonic.*

Proof: Using theorem 8 the map $\tilde{\tau}^\theta$ is \tilde{S}^1 -pluriharmonic. In the special para-Kähler case we know that D is the Levi-Civita connection and hence it is torsion-free. The para-complex structure τ is integrable and hence it holds $N_\tau = 0$. This means, that D is nice and theorem 5 shows that $\tilde{\tau}^\theta$ is para-pluriharmonic. Since S is trace-free we get from proposition 7 that $\tilde{\tau}^\theta$ is harmonic. \square

References

- [CS1] V. Cortés and L. Schäfer, *Topological-antitopological fusion equations, pluriharmonic maps and special Kähler manifolds*, Progress in Mathematics 234, Birkhäuser 2005.

- [CS2] V. Cortés and L. Schäfer, *Flat nearly Kähler manifolds*, submitted.
- [CS3] V. Cortés and L. Schäfer, *A note on flat nearly para-Kähler manifolds*, in preparation.
- [CMMS] V. Cortés, C. Mayer, T. Mohaupt and F. Saueressig, *Special Geometry of Euclidean Supersymmetry I: Vector Multiplets*, J. High Energy Phys. JHEP03(2004)028, hep-th/0312001.
- [CV] S. Cecotti , C. Vafa, *Topological-antitopological fusion*, Nuclear Physics **B 367** (1991), 351-461.
- [D] B. Dubrovin, *Geometry and integrability of topological-antitopological fusion*, Commun. Math. Phys. **152** (1993), 539-564.
- [ET] J.-H. Eschenburg and R. Tibuzy, *Associated families of pluriharmonic maps and isotropy*, Manuscr. Math. **95** (1998), no. 3, 295–310.
- [IZ] S. Ivanov, S. Zamkovoy, *Para-Hermitian and Para-Quaternionic manifolds*, Diff. Geom. Appl. **23** (2005), 205–234.
- [KN] S. Kobayashi, K. Nomizu, *Foundations of differential geometry Volume I/II*, Interscience Publishers 1963/1969.
- [LS] M.-A. Lawn and L. Schäfer, *Decompositions of para-complex vector-bundles and affine para-complex immersions*, Res. in Math. vol. **48**, 3/4.
- [McDS] D. Mc Duff, D. Salamon, *Introduction to symplectic topology*, Oxford University Press (1995).
- [S1] L. Schäfer, *tt^* -geometry and pluriharmonic maps*, Ann. Global Anal. Geom. **28**, no. **3** (2005), 285–300.
- [S2] L. Schäfer, *tt^* -bundles in para-complex geometry, special para-Kähler manifolds and para-pluriharmonic maps*, Diff. Geom. Appl. **24** (2006), 60–89.
- [S3] L. Schäfer, *tt^* -geometry on the tangent bundle of an almost complex manifold*, to appear in J. Geom. Phys.